



<http://www.entm.purdue.edu/Entomology/ext/targets/newslett.htm>

August 8, 2003 - No. 21

In This Issue

Insects, Mites, and Nematodes

- Soybean Aphid Update
- Corn Growth Stage and Insect Potential
- Monitor Soybean for Rootworm Beetles Now
- Black Light Trap Catch Report

Bits & Pieces

- Goodbye Ron Blackwell

Weather Update

- Temperature Accumulations

Insects, Mites, and Nematodes

Soybean Aphid Update – (John Obermeyer and Larry Bledsoe) –

- Aphids are increasing in many fields of northern Indiana
- Now is the time to make treatment decisions
- Pathogens may begin to naturally control aphids

Please refer to the last two issues of the *Pest&Crop* concerning this pest. Soybean aphid numbers continue to climb in research fields in northern Indiana. The most dramatic increase has been at the Pinney Purdue Agricultural Center, Porter County, where aphids per plant increased from 145 to over 600.

Because of ample soil moisture and occasional rains, fields are not showing the presence of aphids or their damage. However, research trials conducted in 2001 indicate that significant yield losses can occur when plants are actively flowering and setting pods with hundreds of aphids per plant. These trials also indicate that treatment decisions need to be made soon, as applications made later in August are too late.

Good news! Diseased aphids are beginning to be seen, they are discolored and shrunk. This disease has been known to literally wipeout aphid populations in a few days time. Be certain to be looking for these while sampling.



Soybean leaflet with over 100 aphids

• • P&C • •

Corn Growth Stage and Insect Potential – (John Obermeyer and Larry Bledsoe) –

- Late developing corn may attract moths
- Fall armyworm whorl feeding is quite evident
- European corn borer infestation is subtle until harvest time

Fall armyworm moths are attracted to **whorl stage corn**. So far, only limited infestations have been reported in southern Indiana and Kentucky. Because of this year's extremely late planting, there are several fields throughout the state that are still quite vulnerable to this vicious feeder. Unless one is willing to spot treat those areas infested with this pest with a high-boy, treatment should not be considered unless 60% of the normal yield is expected, 75% of the plants exhibit feeding, and the larvae are present and actively feeding. This caterpillar is difficult to control unless caught early.

European corn borer moths are attracted to actively **pollinating corn**. Moth flights of the second generation have been low statewide in our black light traps. However, we've received a few reports from pest managers that indicate that "windshield splatter" has been heavy in their area at night. Scouting for egg masses and newly hatched larvae in pollinating corn is very difficult and time consuming. Control of second generation corn borer is difficult, two insecticide applications are often required. In other words, treating for this pest is not likely to be economical for yellow dent corn. High value crops, e.g., sweet and seed corn, should be monitored for egg masses and larvae.

More information on each of these pests can be viewed on-line at <<http://www.entm.purdue.edu/entomology/ext/fieldcropsipm/ciro.htm>>.

• • P&C • •

Monitor Soybean for Rootworm Beetles Now - (John Obermeyer and Larry Bledsoe) -

- Western corn rootworm beetles are active in soybean fields
- Sampling soybean fields now for beetle presence and densities could save money next year
- Sampling doesn't require the use of nets or traps, visual inspections can be effective in assessing relative beetle numbers

We are actively sweeping soybean fields to sample for western corn rootworm beetle variant populations in the state. Though we don't have exact counts yet, it is evident that beetle numbers are down from a year ago. Stay tuned to future *Pest&Crop* issues as we report the catches by county.

Knowledge of beetle numbers in soybean helps one to gauge the potential risk of rootworm damage to next year's corn. Few beetles mean low risk, thus little need for rootworm protection next year. Lots of beetles means higher risk, and the insecticide "insurance" will likely pay off. Several producers and agribusiness personnel throughout the state have initiated a sampling program for their soybean fields. Because of the variability of beetle numbers from field to field, those willing to inspect soybean now may reap the benefits next spring. Sampling for rootworm beetle in soybean fields does not require sticky traps or sweep nets, but they do make decision making more accurate. Visual inspections during the morning hours, while walking through the field and carefully observing the upper canopy, should help you reach a management decision. Soybean fields should be visited weekly until early September. Refer to *Pest&Crop* #19, July 25, 2003, for suggested sampling with sticky traps.

• • P&C • •

| Black Light Trap Catch Report (Ron Blackwell) | | | | | | | | | | | | | | | |
|---|-------------------|-----|-----|------|-----|-----|----|------------------|-----|-----|------|-----|-----|----|--|
| County/Cooperator | 7/22/03 - 7/28/03 | | | | | | | 7/29/03 - 8/4/03 | | | | | | | |
| | VC | BCW | ECB | SWCB | CEW | FAW | AW | VC | BCW | ECB | SWCB | CEW | FAW | AW | |
| Dubois/SIPAC | 1 | 0 | 5 | 0 | 0 | 0 | 1 | 0 | 6 | 1 | 1 | 1 | 0 | 9 | |
| Jennings/SEPAC | 3 | 1 | 1 | 0 | 0 | 1 | 10 | 1 | 5 | 2 | 0 | 0 | 0 | 2 | |
| Knox/SWPAC | 1 | 2 | 0 | 0 | 1 | 0 | 2 | 1 | 1 | 1 | 0 | 0 | 0 | 3 | |
| LaPorte/Pinney Ag Center | 2 | 1 | 1 | 0 | 0 | 1 | 7 | 0 | 0 | 26 | 0 | 0 | 0 | 5 | |
| Lawrence/Feldun Ag Center | 0 | 1 | 0 | 0 | 0 | 1 | 5 | 0 | 0 | 2 | 0 | 0 | 2 | 12 | |
| Randolph/Davis Ag Center | 1 | 1 | 6 | 0 | 0 | 0 | 11 | | | | | | | | |
| Tippecanoe/Throckmorton Ag Center | 0 | 0 | 0 | 0 | 0 | 0 | 1 | | | | | | | | |
| Vermillion/Hutson | | | | | | | | | | 51 | | | | | |
| Whitley/NEPAC | 4 | 0 | 7 | 0 | 0 | 0 | 11 | 6 | 0 | 48 | 0 | 0 | 1 | 10 | |
| BCW = Black Cutworm ECB = European Corn Borer SWCB = Southwestern Corn Borer CEW = Corn Earworm AW = Armyworm FAW = Fall Armyworm VC = Variegated Cutworm | | | | | | | | | | | | | | | |

Bits & Pieces

Goodbye Ron Blackwell – (John Obermeyer) –

Ron Blackwell, IPM Survey Specialist, is leaving us after ten years of hunting and counting insects. He will be sorely missed, as he coordinated many valuable insect IPM programs.

It has been on his heart for several years to pursue a career in law enforcement. When a deputy sheriff position came open, he couldn't resist. Should you misbehave while traveling through Clinton County, you might get to "visit" with Ron!

I know that he'll continue to think of us as his patrol car's windshield gets splattered in the summer. He might even try to identify it.



Ron, we wish you the very best!!!



"Honest Officer Blackwell -- I was just trying to identify these insects for my boss -- please don't give me a ticket for blocking traffic!"

Weather Update


Temperatures as of August 6, 2003

GDD(9) = Growing Degree Days from April 16 (9% of Indiana's corn planted), for corn growth and development
 GDD(26) = Growing Degree Days from April 25 (26% of Indiana's corn planted), for corn growth and development
 GDD(50) = Growing Degree Days from April 30 (50% of Indiana's corn planted), for corn growth and development
 GDD(85) = Growing Degree Days from June 4 (85% of Indiana's corn planted), for corn growth and development

| MAP KEY | | | |
|----------|---------|---------|---------|
| Location | | | |
| GDD(9) | GDD(26) | GDD(50) | GDD(85) |

4" Bare Soil
 Temperatures
 8/6/03

| Location | Max. | Min. |
|----------|------|------|
|----------|------|------|



| | | |
|---------|----|----|
| Wanatah | 86 | 69 |
|---------|----|----|

| | | |
|---------|----|----|
| Winamac | 84 | 66 |
|---------|----|----|

| | | |
|----------|----|----|
| Bluffton | 74 | 73 |
|----------|----|----|

| | | |
|------------|----|----|
| W Laf Agro | 79 | 71 |
|------------|----|----|

| | | |
|-------------|----|----|
| Tipton | 83 | 75 |
| Farmland | 77 | 72 |
| Perrysville | 77 | 72 |

| | | |
|---------|----|----|
| Liberty | 81 | 68 |
|---------|----|----|

| | | |
|---------|----|----|
| Oolitic | 84 | 69 |
|---------|----|----|

| | | |
|--------|----|----|
| Dubois | 91 | 71 |
|--------|----|----|

The **Pest Management and Crop Production Newsletter** is produced by the Departments of Agronomy, Botany and Plant Pathology, and Entomology at Purdue University. The Newsletter is published monthly February, March, October, and November. Weekly publication begins the first week of April and continues through mid-September. If there are questions or problems, contact the Extension Entomology Office at (765) 494-8761.

DISCLAIMER

Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer. It is the policy of the Purdue University Cooperative Extension Service, David C. Petritz, Director, that all persons shall have equal opportunity and access to the programs and facilities without regard to race, color, sex, religion, national origin, age, marital status, parental status, sexual orientation, or disability. Purdue University is an Affirmative Action employer.