A POCKET GUIDE FOR GROWERS

# **BENEFICIAL SYRPHID FLIES**

in Indiana High Tunnels



Allison Zablah & Laura L. Ingwell



Extension - Entomology

# **LIFE CYCLE**

Syrphid flies go through complete metamorphosis, meaning they have four distinct developmental stages: egg, larva, pupa, and adult.

Egg: The female lays tiny, white, oval-shaped eggs directly on the plant surface near aphid colonies, often on leaves or stems.

Larva: After hatching, the legless larva begins feeding on aphids or other softbodied pests. This stage

lasts 7-21 days, depending on the temperature. Larva typically go through 3 instars (stages) before pupating, increasing in size and the amount of prey they consume as they grow.

Pupa: When fully grown, the larva drops to the soil or stays on the plant to form a hard case (pupa), inside of which the final transformation to adult occurs. This stage lasts 7-14 days, depending on temperature, but can be shorter in hot weather or longer in cooler conditions.

Adult: The adult fly emerges from the pupal case. Males and females look very similar within a species. Both feed on nectar and pollen. contributing to pollination in the process. Adults are sexually mature in 2-5 days, then mate and reproduce. Multiple generations can occur during the growing season, especially in warm weather.

# INTRODUCTION

Syrphid flies, also known as hoverflies or flower flies, are beneficial insects commonly mistaken for bees. However, they are true flies and can be distinguished by having only one pair of wings, unlike bees, which have two pairs (four wings). As adults, syrphids play a valuable role as pollinators, while their larvae serve different functions depending on the species. They can be predators,

feeding on soft-bodied pests, pollen feeders, like aphids, or decomposers, helping break down decaying organic matter. This pocket guide highlights the most common aphid-eating (aphidophagous) syrphid species found in high tunnel systems, with images and descriptions Photos by John Obermeyer (left) and of both the adult and larval stages.



Allison J. Zablah (right: larval stages)

Why do syrphids matter in agriculture? Learning to identify and recognize these helpful insects can lead to more informed pest management decisions, potentially reducing pesticide use and allowing natural enemies to manage pest populations. The syrphids featured in this guide are found specifically in high tunnel environments, a habitat that is stressful and avoided by many natural enemies. Interestingly, syrphids are also one of the first natural enemies found in spring and last to be seen in fall, highlighting their ability to forage into the shoulder seasons under protected culture.

## **IDENTIFICATION GUIDE**

Adults: Syrphid flies often mimic bees or wasps in shape and color to avoid predators. Most have slender or oval-shaped bodies and range in size from 4 mm to over 15 mm, depending on the species. Despite looking like bees or wasps, they do not sting or bite. Below are some key features:



Photo by Allison J. Zablah

Larvae: Syrphid fly larvae are maggots: legless with a tapered body. They are commonly green, brown, or translucent, making them well camouflaged on plant surfaces. You'll frequently find them in or around aphid colonies, especially on the underside of leaves or stems of plants. Although they move slowly, these larvae are active predators feasting on soft-bodied prev.



Photos by Allison J. Zablah

# **HOW TO SUPPORT SYRPHIDS ON YOUR FARM**

## 1. Provide Continuous Floral Resources

- Plant flowering plants that bloom at different times to offer nectar and pollen throughout the season. Best options include:
  - ▶ Early-season: sweet alyssum, cilantro, and buckwheat.
  - ▶ Mid-season: dill, fennel, coriander, and mustard greens.
  - Late-season: sunflowers, goldenrod, and
- Plant flowers in strips within or next to crops, or simply add flowering plants along the outer edges of the field to attract beneficial insects.

## 2. Reduce Soil Disturbance

- Syrphid fly larvae often drop to the soil to
- Minimize tillage to avoid killing pupae.

#### 3. Minimize Pesticide Use

- Syrphid flies are susceptible to insecticides.
- Use Integrated Pest Management (IPM) strategies:
  - ▶ Monitor pest levels.
  - ▶ Use thresholds before spraying.
  - Apply selective or reduced-risk products in the evening when flies are less active.
  - Avoid contact insecticides like pyrethroids and organophosphates.

# 4. Maintain Habitat Diversity

- Provide a mix of crops and wild plants; diversified landscapes attract more beneficial insects.
- Maintain hedgerows, grassy margins, and native vegetation nearby providing perennial habitat.

# 5. Provide Shelter and Overwintering Sites

- Adult syrphids overwinter in leaf litter, grassy areas, or undisturbed field margins.
- Avoid burning or clearing these habitats in fall
- Wait to mow cover crops or wildflower strips until syrphid flies are active in spring, so you don't destroy overwintering adults.

## **ACKNOWLEDGMENTS**

The life cycle illustration was created by Sheyla Zablah and the abdomen illustrations were created by Allison Zablah. Guide designed by Bransen Shidler.

This work is supported by the Specialty Crop Research Initiative (SCRI) [Grant No. 2021-51181-35858 / Project Accession No. 1027430] from the USDA National Institute of Food and Agriculture.



USDA National Institute of Food and Agriculture U.S. DEPARTMENT OF AGRICULTURE

# **HOW TO USE THIS GUIDE AND WHY IT MATTERS**


This pocket guide is designed as a practical tool to help you identify syrphid flies (also known as hoverflies) commonly found in Indiana high tunnel crop systems. Carry it with you to recognize beneficial syrphid species on the spot. Understanding which syrphids are present can give you insight into your local insect community and help you improve pest management strategies, especially for controlling aphids and other soft-bodied pests. Many syrphid larvae are natural predators of aphids, while adults feed on nectar and pollen, contributing to pollination. Syrphid larvae are often hard to spot because they blend in with the plant surface, so recognizing the adults is the first step to scouting for this beneficial insect. If you see adults present, check aphid, thrips, or mite populations closely, as these pests may already be under natural control due to syrphid larvae feeding on them. This is especially important in protected environments like high tunnels, where traditional pollinators like bees are not abundant. Soft-bodied pests such as aphids, thrips, and mites are common in these systems, providing abundant food for syrphid larvae. By learning to recognize syrphids, you can better support and conserve these beneficial insects, contributing to healthier crops and reduced pesticide use. Know your bugs because the right ID can lead to more resilient farms!


# **OTHER TOOLS & RESOURCES**


- If you'd like to learn more about syrphid flies, we recommend the book Field Guide to the Flower Flies of Northeastern North
- For information on specialty crop research conducted in high tunnels, scan the QR code
- If you're looking for identification keys, see Flies of Illinois, USA, page 8.
- Online key to genera (Skevington et al., p. 19)
- You can also visit Purdue's Vegetable Crops Hotline, particularly Issue #751 and Issue #737, for more practical information.


# REFERENCES

- Skevington, J. H., Crins, W. J., Locke, M. M., Marshall, S. A., & Young, A. D. (2019). Field Guide to the Flower Flies of Northeastern North America, Princeton University Press. https://doi.org/10.1515/9780691192512
- Dunn, L., Lequerica, M., Reid, C. R., & Latty, T. (2020). Dual ecosystem services of syrphid flies (Diptera: Syrphidae): pollinators and biological control agents. Pest Management Science, 76(6), 1973-1979. https://doi.org/10.1002/ps.5807
- Rotheray, G. E., & Gilbert, F. (2022). Syrphid (Diptera: Syrphidae) biology: Evolution, ecology, and ecosystem services. Insects, 13(7), 573. https://doi.org/10.3390/
- Miranda, G. F. G., Young, A. D., Locke, M. M., Marshall, S. A., Skevington, J. H., & Thompson, F. C. (2013). Key to the genera of Nearctic Syrphidae. Canadian Journal of Arthropod Identification, (23), https://doi.org/10.3752/ciai,2013.23
- Illinois Nature Preserves Commission, & Moorehouse, A. (2020), Flies of Illinois (Guide No. 1255, 16 pp.), Field Museum of Natural History.
- Davis, A. E., Bickel, D. J., Saunders, M. E., & Rader, R. (2023). Crop-pollinating Diptera have diverse diets and habitat needs in both larval and adult stages. Ecological Applications, 33(5), e2859. https://doi.org/10.1002/eap.2859












Photo by Steven Falk, Flickr. CC BY 2.0 (https://creativecommons.org/licenses/by/2.0/)

